



Shahrood University of
Technology

Journal of Mining and Environment (JME)

Journal homepage: www.jme.shahroodut.ac.ir



Iranian Society of
Mining Engineering
(IRSM)

Dissolution Kinetics of Metallic Silver using Ammonium Nitrate and Hydrogen Peroxide Solution as an Environmentally Friendly Cyanide Substitute

Salih Aydogan¹, Mohamed Taha Osman Abdelraheem^{1,2*}, Babiker Ali³, and Mustafa Boyrazli⁴

1. Department of Mining Engineering, Faculty of Engineering and Natural Sciences, Konya Technical University, Konya, Turkey

2. Department of Mining Engineering, Faculty of Engineering Sciences, Omdurman Islamic University, Omdurman, Sudan

3. Department of Economic Geology and Mining, Red Sea University, Port Sudan, Sudan

4. Department of Metallurgical and Materials Engineering, Faculty of Engineering, Firat University, Elazig Turkey

Article Info

Received 8 April 2025

Received in Revised form 28 May 2025

Accepted 15 September 2025

Published online 15 September 2025

DOI: [10.22044/jme.2025.16031.3089](https://doi.org/10.22044/jme.2025.16031.3089)

Keywords

Ammonium nitrate

Hydrogen peroxide

Dissolution

Silver

Activation energy

Abstract

This article describes the kinetics of utilizing ammonium nitrate to dissolve pure metallic silver in hydrogen peroxide solution (H₂O₂). Using pure metallic silver allows for precise leaching kinetics research by removing interference from impurities and facilitating accurate interpretation of rate-controlling mechanisms. The impact of temperature, rotation speed, H₂O₂ concentration, and ammonium nitrate concentration were all examined. The results show a favourable relationship between the rate of silver (Ag) dissolution and the rotation speed. Additionally, a low concentration of ammonium nitrate (between 0.003 and 0.20 M) has advantageous effects on Ag dissolution. The dissolution rate was significantly impacted by H₂O₂ concentrations between 0.08 and 0.15 M, because this range of H₂O₂ concentration required to provide sufficient oxidative potential for significant silver solubility. However, this effect is less pronounced in the 0.20–0.50 M range. 20 - 50 °C range of temperatures are advantageous since H₂O₂ is stable in this range. It was calculated that the activation energy was 25.66 kJ/mol.

1. Introduction

Silver (Ag) is an essential metal that is used in jewelry, dental amalgam, electrical and electronic equipment, photographic materials, catalysts, and brazing alloys [1]. The cyanidation process is the most widely used extraction method for gold and silver extraction at the moment [2].

Cyanide is highly toxic and can harm both human health and the environment, despite its many benefits [3]. In order to identify substitutes that are less detrimental to the environment and public health, considerable investigations have been conducted [4]. Many hydrometallurgical methods have recently been developed to recover silver without the use of cyanide. Many studies have concentrated on the

thiourea [5, 6], thiocyanate [7-9], halides [10–12], and thiosulfate techniques [13–16].

Thiourea (NH₂CSNH₂) is a less hazardous chemical for both humans and the environment, which is the reason it was selected to be researched as one of the cyanide substitutes. When using thiourea leaching, acidic conditions are usually employed. When using acidic thiourea as a leaching reagent, a number of problems occur such as high thiourea consumption when compared to cyanide, challenging thiourea lixiviant regeneration and substantial equipment corrosion in acidic solutions compared to steel passivation in alkaline cyanide solutions [17]. These problems



Corresponding author: mohamedtaha@oiu.edu.sd (M.T.O. Abdelraheem)

have prevented thiourea from being used in industry.

Thiocyanate (SCN^-) is also used as an alternative for cyanide to extract silver and gold. Thiocyanate produced when sulfur species react with cyanide. Thiocyanate solutions are less toxic and more stable than potassium and sodium cyanide. It is also renewable and accessible as a result of technological processes.

Moreover, it has been found that the halides are efficient agents for leaching gold and silver in water-based solutions. Typically, the halogen species functions as both a ligand and an oxidant in halide processes. The pH, solution potential, presence of reductants, and halide concentration all have a substantial impact on the stability of the halide complexes during the leaching process with halide ions [18]. For the halide systems to reach the high oxidation potentials needed for leaching, an oxidant such as bromine (Br_2) or iodine (I_2) must be used, but both are toxic [19].

Thiosulfate ($\text{S}_2\text{O}_3^{2-}$) is another alternative to cyanide that is less dangerous. Up to 12 g of thiosulfate can be taken orally every day without risk [19]. Thiosulfate leaching has gained favor over cyanide leaching because it is a non-toxic reagent and leaches more quickly. But the main problems with the thiosulfate leaching system are that it uses a lot of reagents and does not have a good method for recovering gold [20, 21].

Despite considerable efforts to develop environmentally benign alternatives to traditional cyanide-based leaching reagents for silver extraction, these substitutes still present several limitations. Common drawbacks include high operational costs, limited potential for reagent recycling, and challenges in maintaining control over key process variables. These issues underscore the need for continued research into effective cyanide alternatives.

Although Starovoytov [22] investigated the dissolution of silver (Ag) in ammonia solution, Aydogan et al. [23,24] studied the dissolution of Ag in ammonium salts, noting that ammonia is volatile and can easily escape from open leaching vessels, potentially causing environmental harm. Aydogan et al. [23,24] concluded that using only ammonia solution as a leaching agent for silver extraction is not advisable. Instead, they recommended the use of ammonium salts such as ammonium carbonate, diammonium tartrate, or ammonium nitrate.

Since the leaching kinetics of ammonium carbonate and diammonium tartrate have already been studied, the authors chose to investigate

ammonium nitrate in this study to evaluate the feasibility of using an ammonium nitrate and hydrogen peroxide solution as an alternative leaching system for metallic silver.

Ammonium nitrate, commonly used as a chemical fertilizer in agriculture, known for its high solubility and availability [25,26], making it an attractive candidate for hydrometallurgical applications. When combined with hydrogen peroxide, the system can provide the oxidative potential required to dissolve metallic silver while minimizing environmental impact. This approach avoids the generation of harmful byproducts, aligning well with the principles of green chemistry. The work specifically aims to investigate the leaching kinetics of this eco-friendly system under various experimental conditions, including changes in temperature, reagent concentration, and reaction time. By analyzing these variables, the study aims to determine the reaction order, activation energy, and potential rate-controlling steps.

2. Materials and Methods

A high-purity metallic silver disc (99.99%) was used in all leaching experiments. The disc had a surface area of 23.85 cm^2 and a thickness of 3 mm. Ammonium nitrate (NH_4NO_3), used as the leaching reagent, was obtained from Merck, as was the hydrogen peroxide (H_2O_2) solution employed as the oxidizing agent.

A silver disc was attached to a Teflon-coated mechanical shaft, which was immersed in a 1000-milliliter beaker containing the leaching solution and placed in a temperature-controlled water bath for the experiments (Figure 1). The effects of 0.08–0.50 M H_2O_2 , rotating speed between 314 rad/min and 942 rad/min, 0.03–0.20 M ammonium nitrate, and 20–50 °C temperature ranges were investigated. In each test, half a liter of solution was used to extract the metallic silver over a period of 24 min. The dissolved silver was analyzed with an atomic absorption spectrophotometer (GBC SensAA model). Samples of the leaching solution were taken frequently. High-quality reagents and distilled water were used for each experiment.

3. Results and Discussion

3.1. Leaching Mechanism of Ag with ammonium nitrate

To comprehend the mechanism of formation of ammonia complex with Ag, it is imperative to examine the impact of pH on Ag dissolution. The experiment was run with the following parameters:

30 °C, 0.15 M ammonium nitrate concentration, 314 rad/min rotation speed, and 0.15 M hydrogen peroxide concentration. The pH value under these circumstances was found to be 4.84. NaOH was used to raise the pH above 4.84. As shown in Figure 2, the solubility of silver increases in an alkaline medium with increasing pH. This experiment indicates that ammonia and silver combine to form a stable complex under conditions of an alkaline




Figure 1. Experimental set-up.

In this situation, the following equations can be used to explain the mechanism of Ag dissolution in alkaline medium. Equation 2 demonstrates that H_2O_2 has a strong oxidizing agent value of 1.007 V in an alkaline medium [27].



The following formulas can be used to understand the mechanism of silver-ammonia complex formation [27]:





Figure 3 also demonstrates that silver is soluble in acidic media. The reaction orders with respect to solution pH were determined to be 0.009 in acidic medium and 0.17 in alkaline medium.

pH. The stability of the ammonia-silver complex is demonstrated by equation (1), which also shows a relatively high formation constant ($K_f = 1.1 \times 10^7$) [27].







Figure 2. Impact of pH on the leaching of Ag rotating disc in alkaline medium (conditions: rotation speed of 314 rad/min, 0.15 M ammonium nitrate concentration, 30 °C, 23.85 cm² disc surface area and 0.15 M H_2O_2).



Equation 5 and 6, which expresses the final equations describing the dissolution of Ag with NH_4NO_3 in an alkaline medium, can be derived from Equations 3 and 4.

As shown in Figure 4, the reaction order obtained for the acidic medium is lower than that for the alkaline medium, indicating that silver dissolution is minimally affected by changes in pH under acidic conditions.




Figure 3. Impact of pH on the leaching of Ag rotating disc in acidic medium (conditions: rotation speed of 314 rad/min, 0.15 M ammonium nitrate concentration, 30 °C, 23.85 cm² disc surface area and 0.15 M H₂O₂).




Figure 4. The reaction orders in relation to pH (conditions: rotation speed of 314 rad/min, 0.15 M ammonium nitrate concentration, 30 °C, 23.85 cm² disc surface area and 0.15 M H₂O₂).





Figure 5. The Eh-pH diagram of silver-ammonia-water (Ag-NH₃-H₂O) system [23]


The Pourbaix diagram of the NH₃-H₂O-Ag system (Figure 5) suggests that an acidic environment inhibits the formation of silver-

ammonia complexes. In this case, the following formulas can be used to explain the mechanism of silver dissolution.

Equation 7 shows that H₂O₂ has a 1.763 V redox potential value in acidic condition [24].



The following formulas can be used to explain the mechanism of silver nitrate formation [24]:



Equation 10 and 11, can be utilized to express the final equations that characterizes the leaching of Ag with NH₄NO₃ in an acidic medium, based on Equations 8 and 9.



3.2. Effect of Ammonium Nitrate Concentration

At a 314 rad/min rotation speed, a temperature of 30 °C, and a concentration of 0.15 M H₂O₂, the effect of ammonium nitrate concentration on Ag dissolution is shown in Figure 6(a). The dissolution of Ag was found to be significantly influenced by the concentration of ammonium nitrate, which ranges from 0.003 to 0.200 M. Increasing concentration of ammonium nitrate causes a proportion of free ammonia to rise, which facilitates the formation of ammonia- silver

complexes and explains why the dissolution of Ag is increasing.

A logarithmic linearization method was used to determine the reaction order in order to quantify the impact of ammonium nitrate. The natural logarithm of the dissolution rate (ln(rate)) was plotted against the natural logarithm of the ammonium nitrate concentration (ln [NH₄NO₃]). The resulting linear plot, shown in Figure 6(b), yielded a slope of 0.66, indicating a reaction order less than unity. This

result suggests that while the dissolution rate of silver increases with ammonium nitrate

concentration, the relationship is not directly proportional.




Figure 6. (a) Influence of ammonium nitrate concentration on the Ag leaching (conditions: rotation speed of 314 rad/min, 30 °C, 23.85 cm² disc surface area and 0.15 M H₂O₂); (b) The reaction order in relation to ammonium nitrate concentration.

3.3. Effect of H₂O₂ Concentration

Using a 0.15 M ammonium nitrate concentration, 314 rad/min rotation speed, and 30 °C temperature, Figure 7(a) shows the Ag dissolution versus H₂O₂ concentration in the 0.080–0.50 M range. The results show that silver dissolves more readily when H₂O₂ concentration is raised to a range of 0.08–0.15 M. However, at a concentration of 0.20–0.50 M, there is no appreciable changes on the Ag dissolution. The chemical reaction order of H₂O₂ concentration in silver dissolution was determined to be 0.80 and for the concentrations of 0.08–0.15 M and 0.20–0.50 M, respectively, based on the linearized plots in Figure 7(b).

The concentrations of H₂O₂ exhibit two slopes, as illustrated in Figure 7(b). This behavior can be attributed to the role of H₂O₂ as an oxidizing agent, which facilitates the reaction between silver and ammonium nitrate. The concentration range of H₂O₂ required to provide sufficient oxidative potential for significant silver solubility is between 0.08 and 0.15 M. Beyond this range, particularly at concentrations above 0.20 M, the slope of the curve approaches zero (0.07), indicating that further increases in H₂O₂ concentration have a negligible effect on silver dissolution. Aydoğan et al. [27]

noted that at high H₂O₂ concentrations, an oxygen layer formed on the silver disc.

3.4. Impact of Temperature

The impact of temperature between 20–50 °C on the leaching rate of Ag at a rotation speed of 314 rad/min, an ammonium nitrate concentration of 0.15 M, and a H₂O₂ concentration of 0.15 M presented in Figure 8(a). It demonstrates how rising temperatures cause silver to dissolve more readily. Since hydrogen peroxide normally breaks down at temperatures above 60 °C, the rate at which silver dissolves has not been investigated at higher temperatures [28, 29].

After calculation based on the Arrhenius plot, the activation energy of the process was found to be 25.66 kJ/mol as illustrated in Figure 8(b). This value falls within the typical range for mixed kinetic control (12–39 kJ/mol) [30, 31] suggesting that the leaching process is controlled by a combination of surface chemical reaction and diffusion mechanisms. Similar observations have been reported in the literature. For instance, Yang and Honaker [32] reported an activation energy of 27 kJ/mol, which indicated a mixed control mechanism. Likewise, Xu et al. [33] obtained a value of 22.056 kJ/mol in their study on willemite leaching, also supporting the interpretation of mixed kinetic behavior.




Figure 7. (a) Influence of H_2O_2 concentration on the Ag leaching (conditions: 314 rad/min rotation speed, 30 °C, ammonium nitrate concentration of 0.15 M, 23.85 cm² disc surface area); (b) reaction order regarding to H_2O_2 concentration




Figure 8. (a) Impact of temperature on the leaching of Ag (conditions: ammonium nitrate concentration of 0.15 M, rotation speed of 314 rad/min, 0.15 M H_2O_2 , 23.85 cm² disc surface area); (b) reaction order in relation to temperature

3.5. Impact of Rotating Speed

The impact of rotation speed on the dissolution of silver at an ammonium nitrate of 0.15 M, a temperature of 30 °C, and a H_2O_2 concentration of 0.15 M presented in Figure 9. The results proved that the rotation speed has a good impact on the dissolution rate of Ag, which has an order of 0.388, as can be seen in Figure 9(b). Higher rotation speeds enhance the leaching rate by improving mass transfer at the solid–liquid interface.

Additionally, the increase in the rotation speeds ensures a more homogeneous distribution of oxidants and complexing agents, which leads to more consistent dissolution rates. These results are in agreement with previous studies, such as Aydogan et al. [27], who reported that the silver leaching rate was significantly influenced by the rotation speed. Based on the supposition that there is laminar flow on the surface of the rotating disc [34], the Levich equation is validated by the result from Figure 10.




Figure 9. (a) Impact of rotation speed on the Ag leaching (conditions: 0.15 M H_2O_2 , ammonium nitrate concentration of 0.15 M, 30 °C, disc surface area of 23.85 cm^2); (b) reaction order according to rotation speed.

3.6. Impact of Rotating disc surface area

Figure 11 (a) shows the impact of the disc surface area on the Ag leaching rate at an ammonium nitrate concentration of 0.15 M, a temperature of 30 °C, a rotation speed of 314 rad/min, and a H_2O_2 concentration of 0.15 M. Increases in surface area have been found to accelerate the rate at which silver dissolves. As shown in Figure 11(b), the slope of approximately 1. In this case, Levich and Tobias assume that the dissolution rate of Ag is directly proportional to the surface of disc [34].




Figure 10. The relationship between the leaching rates and $\omega^{1/2}$ at 0.15 M H_2O_2 , ammonium nitrate concentration of 0.15 M, 30 °C, 23.85 cm^2 disc surface area).




Figure 11. (a) Impact of disc surface area on the Ag leaching (conditions: 0.15 M H_2O_2 , ammonium nitrate concentration of 0.15 M, 30 °C, rotation speed of 314 rad/min); (b) reaction order according to disc surface area

3.7. Derivation of the Overall Kinetic Model

In 1963, Levich and Tobias presented an equation for determining the rate of dissolution of the rotating disc [34]. The amount of dissolved metal is established by this equation as a function

$$X = f(\omega, A, C_{\text{hydrogen peroxide}}, C_{\text{ammonium nitrate}}, t, \exp(-E/RT)) \quad (12)$$

Where are:

A - surface area (cm²)

E - activation energy (kJ/mol),

k - rate constant

X - mass of silver dissolved (mg)

ω - the rotational speed of the disc (rad/min)

of multiple parameters—such as rotation speed, surface area, concentrations of hydrogen peroxide and ammonium nitrate, time, and temperature—as expressed in Equation (12). This comprehensive approach makes it more robust and informative compared to many previous models.

$C_{\text{ammonium nitrate}}$ - the concentration of ammonium nitrate(M)

and $C_{\text{hydrogen peroxide}}$ - the hydrogen peroxide concentration (M)

R - the universal gas constant

By changing the parameters, equation (13) was produced.

$$X = k \omega^e A^f C_{\text{hydrogen peroxide}}^g C_{\text{ammonium nitrate}}^h e^{-E/RT} t \quad (13)$$

where e, f, g, and h are constants to be found in this work.

The following empirical model can be used to express the degree of the Ag dissolution (X) of this process:

$$X = 4.419 \times 10^3 \omega^{0.5} A^{0.999} C_{\text{hydrogen peroxide}}^{0.80} C_{\text{ammonium nitrate}}^{0.66} e^{-25660/RT} t \quad (14)$$




Figure 12. Empirical vs. experimental dissolved Ag

Ammonium nitrate has an order (0.66) less than that of hydrogen peroxide (0.80). These orders indicate that the most effective parameter in the system was the hydrogen peroxide.

As demonstrated by the linear correlation in Figure 12 with a slope of 1 and a high coefficient of determination ($R^2 = 0.996$), this model exhibits

strong agreement with the experimental results, confirming the predictive accuracy of the model.

4. Conclusions

This work examined the leaching kinetics of pure Ag using ammonium nitrate in H₂O₂ solution instead of cyanidation process due to the environmental problems of cyanide. The dissolution kinetics of Ag were studied by investigating the influences of temperature, ammonium nitrate concentration, H₂O₂ concentration, and rotating speed on the dissolution rate. The findings show that a small concentration of ammonium nitrate (0.15 M) was sufficient to dissolve silver alternatively to cyanide. This suggests that ammonium nitrate may serve as a viable, potentially safer alternative to cyanide in Ag extraction processes. It was found that the H₂O₂ concentrations between 0.08-0.15 M are enough to provide adequate oxidative potential to achieve a high degree of Ag solubility. Increasing the H₂O₂ concentration beyond 0.20 M did not result in further improvement in dissolution efficiency, indicating a saturation point in oxidative activity.

Kinetic analysis yielded an apparent activation energy of 25.66 kJ/mol

The study recommends further application of ammonium nitrate leaching to silver-bearing ores and comparative analysis with traditional cyanidation methods to evaluate industrial feasibility. Such investigations will help establish the practicality and environmental benefits of adopting ammonium nitrate-based leaching systems for large-scale silver extraction.

Acknowledgments

The authors would also like to thank the Konya Technical University for guidance and providing information.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- [1]. Aydogan, S., Abdelraheem, M. T. O., Ali, B., & Boyrazli, M. (2025). Kinetics Study on the Leaching of Metallic Silver with Ammonium Carbonate as an Eco-Friendly Alternative for Cyanide. *JOM*, 1-10.
- [2]. Osman Abdelraheem, M. T., Akasha, M., & Agacayak, T. (2022). Recyanidation of Gold Heaps Tailing at Hassai Region in Red Sea State, Sudan. *Journal of Mining Science*, 58(4), 628-634.
- [3]. Chen, X., Ren, Y., Qu, G., Wang, Z., Yang, Y., & Ning, P. (2023). A review of environmental functional materials for cyanide removal by adsorption and catalysis. *Inorganic Chemistry Communications*, 157, 111298.
- [4]. Islas, H., Flores, M. U., Juárez, J. C., Reyes, M., Blanco, A., Gutierrez, E. J., ... & Reyes, I. A. (2021). Silver leaching from jarosite-type compounds using cyanide and non-cyanide lixiviants: A kinetic approach. *Minerals Engineering*, 174, 107250.
- [5]. Abikak, Y., Kenzhaliev, B., Akcil, A., Dembele, S., Koizhanova, A., Bakhtuly, N., & Kassymova, G. (2025). Optimization of Thiourea-Promoted Gold and Silver Leaching from Pyrite Cinders Using Response Surface Methodology (RSM). *Processes*, 13(5), 1277.
- [6]. Gamiño-Arroyo, Z., Pareau, D., Buch, A., Gomez-Castro, F. I., Sánchez-Cadena, L. E., Stambouli, M., ... & Avila-Rodriguez, M. (2021). Design of multistage extraction system for simultaneous separation of silver and gold from thiourea solutions. *Chemical Engineering and Processing-Process Intensification*, 164, 108391.
- [7]. Azizitorghabeh, A., Wang, J., Ramsay, J. A., & Ghahreman, A. (2021). A review of thiocyanate gold leaching—Chemistry, thermodynamics, kinetics and processing. *Minerals Engineering*, 160, 106689.
- [8]. Rezaee, M., Abdollahi, H., Saneie, R., Mohammadzadeh, A., Rezaei, A., Darvanjooghi, M. H. K., ... & Magdoui, S. (2022). A cleaner approach for high-efficiency regeneration of base and precious metals from waste printed circuit boards through stepwise oxido-acidic and thiocyanate leaching. *Chemosphere*, 298, 134283.
- [9]. Azizitorghabeh, A., Mahandra, H., Ramsay, J., & Ghahreman, A. (2021). Gold leaching from an oxide ore using thiocyanate as a lixiviant: process optimization and kinetics. *ACS omega*, 6(27), 17183-17193.
- [10]. El-Shaheny, R., El Hamd, M. A., El-Enany, N., Alshehri, S., & El-Maghrabey, M. (2024). Insights on the utility of ionic liquids for greener recovery of gold and silver from water, wastes, and ores. *Journal of Molecular Liquids*, 126034.
- [11]. Vereycken, W., Riano, S., Gerven, T. V., & Binnemans, K. (2020). Extraction behavior and separation of precious and base metals from chloride, bromide, and iodide media using undiluted halide ionic liquids. *ACS Sustainable Chemistry & Engineering*, 8(22), 8223-8234.
- [12]. Pereira, M. M., Costa, F. O., Gomes, R. F., Rodrigues, M. L. M., da Silva, G. A., & Leão, V. A. (2020). Multivariate study of a novel hydrometallurgical route employing chloride/hypochlorite for leaching silver from printed circuit boards. *Chemical Engineering Research and Design*, 163, 115-124.
- [13]. Chen, J., Xie, F., Wang, W., Fu, Y., & Wang, J. (2022). Leaching of gold and silver from a complex sulfide concentrate in copper-tartrate-thiosulfate solutions. *Metals*, 12(7), 1152.
- [14]. Hernández-Ávila, J., Salinas-Maldonado, R. G., García-Cerón, A., Flores-Badillo, J., Cerecedo-Sáenz, E., Toro, N., & Salinas-Rodríguez, E. (2025). A Comparative Study of Cyanide and Thiosulfate for Silver Leaching from Tailings: A Kinetics Approach. *Processes*, 13(5), 1522.
- [15]. Roldán-Contreras, E., Salinas-Rodríguez, E., Hernández-Ávila, J., Cerecedo-Sáenz, E., Rodríguez-Lugo, V., I. Jeldres, R., & Toro, N. (2020). Leaching of silver and gold contained in a sedimentary ore, using sodium thiosulfate; a preliminary kinetic study. *Metals*, 10(2), 159.
- [16]. Bae, M., Kim, S., Sohn, J., Yang, D., & Lee, H. (2020). Leaching behavior of gold and silver from concentrated sulfide ore using ammonium thiosulfate. *Metals*, 10(8), 1029.
- [17]. Liu, Z. W., Guo, X. Y., Tian, Q. H., & Zhang, L. (2022). A systematic review of gold extraction: Fundamentals, advancements, and challenges toward alternative lixiviants. *Journal of Hazardous Materials*, 440, 129778.
- [18]. Whitehead, J. A., Zhang, J., McCluskey, A., & Lawrance, G. A. (2009). Comparative leaching of a

sulfidic gold ore in ionic liquid and aqueous acid with thiourea and halides using Fe (III) or HSO_5^- oxidant. *Hydrometallurgy*, 98(3-4), 276-280.

[19]. Lewis, R. J., 1997, "Hazardous chemicals desk reference", New York, international Thomson publishing company.

[20]. Molleman, E. and Dreisinger, D., 2002, "The treatment of copper-gold ores by ammonium thiosulfate leaching", *Hydrometallurgy*, 66, 1-21.

[21]. Navarro, P., Vargas, C., Villarroel, A. and Alguacil, F. J., 2002, "On the use of ammoniacal/ammonium thiosulphate for gold extraction from a concentrate", *Hydrometallurgy*, 65, 37-42.

[22]. Starovoytov, O. N., Kim, N. S., & Han, K. N. (2007). Dissolution behavior of silver in ammoniacal solutions using bromine, iodine and hydrogen-peroxide as oxidants. *Hydrometallurgy*, 86(1-2), 114-119.

[23] Aydogan, S., Abdelraheem, M. T. O., Ali, B., & Boyrazli, M. (2025). Dissolution of metallic silver with ammonium acetate and hydrogen peroxide solution as a greener substitute for cyanide: kinetics study. *Discover Chemistry*, 2(1), 196.

[24]. Aydogan S, A

bdelraheem M T O, Ali B, & Boyrazli M. (2024). Leaching Kinetics of Metallic Silver Using Diammonium Tartrate and Hydrogen Peroxide Solution as a Greener Substitute for Cyanide. *Journal of Sustainable Metallurgy*, 10(4), 2442-2454.

[25]. Yahaya, S. M., Mahmud, A. A., Abdullahi, M., & Haruna, A. (2023). Recent advances in the chemistry of nitrogen, phosphorus and potassium as fertilizers in soil: A review. *Pedosphere*, 33(3), 385-406.

[26]. Kudryashova, O. S., Kataev, A. V., & Malinina, L. N. (2015). Solubility in the NaNO_3 - NH_4NO_3 - KNO_3 -

H_2O system. *Russian Journal of Inorganic Chemistry*, 60, 355-361.

[27]. Aydogan S, Abdelraheem M T O, Ali B, & Boyrazli M. (2025). Leaching Kinetics of Metallic Silver with Sodium Cyanide in Hydrogen Peroxide Solution. *The Canadian Journal of Chemical Engineering*.

[28]. Abdelraheem M T O, Agacayak T (2023) Investigation of the effect of some polar organic solvents on the leaching and dissolution kinetics of chalcopryrite in hydrogen peroxide and sulfuric acid solution. *Bulletin of the Chemical Society of Ethiopia*. 37(3): 779-788.

[29]. Abdelraheem M T O, Agacayak T (2022). Effect of organic and inorganic compounds on dissolution kinetics of chalcopryrite in hydrogen peroxide-Hydrochloric acid system. *Journal of Saudi Chemical Society*, 26(3).

[30]. Feng X, Long Z, Cui D, Wang L, Huang X, Zhang G (2013) Kinetics of rare earth leaching from roasted ore of bastnaesite with sulfuric acid. *Trans. Nonferrous Metals Soc. China* 23: 849-854.

[31]. Saxena N N, Mandre N R (1992) Mixed control kinetics of copper dissolution for copper ore using ferric chloride. *Hydrometallurgy*. 28: 111-117.

[32]. Yang, X. & Honaker, R. Q. (2020). Leaching kinetics of rare earth elements from fire clay seam coal. *Minerals*, 10(6), 491.

[33]. Xu, H., Qian, Y., Zhou, Q., Wei, C., Wang, Q., Zhao, W., ... & Xu, J. (2022). Kinetics and Mechanisms of Artificial Willemite Leaching in Low-Sulfuric-Acid Solution at Elevated Temperature. *Metals*, 12(12), 2031.

[34]. Levich V G, Tobias C W (1963) Physicochemical hydrodynamics. *J. Electrochem. Soc.* 110, 251C.



دانشگاه صنعتی شاهرود

نشریه مهندسی معدن و محیط زیست

www.jme.shahroodut.ac.ir نشانی نشریه:



انجمن مهندسی معدن ایران

سینتیک انحلال نقره فلزی با استفاده از محلول نیترات آمونیوم و پراکسید هیدروژن به عنوان جایگزین سیانید سازگار با محیط زیست

صالح آیدوغان^۱، محمد طه عثمان عبدالرحیم^{۱،۲*}، بابیکر علی^۳، و مصطفی بویرازلی^۴

۱. گروه مهندسی معدن، دانشکده مهندسی و علوم طبیعی، دانشگاه فنی قونیه، قونیه، ترکیه

۲. گروه مهندسی معدن، دانشکده علوم مهندسی، دانشگاه اسلامی ام‌دورمان، ام‌دورمان، سودان

۳. گروه زمین‌شناسی اقتصادی و معدن، دانشگاه دریای سرخ، بندر سودان، سودان

۴. گروه مهندسی متالورژی و مواد، دانشکده مهندسی، دانشگاه فرات، الازیغ ترکیه

چکیده

این مقاله سینتیک استفاده از نیترات آمونیوم برای حل کردن نقره فلزی خالص در محلول پراکسید هیدروژن (H_2O_2) را شرح می‌دهد. استفاده از نقره فلزی خالص با حذف تداخل ناشی از ناخالصی‌ها و تسهیل تفسیر دقیق مکانیسم‌های کنترل سرعت، امکان تحقیق دقیق سینتیک لیچینگ را فراهم می‌کند. تأثیر دما، سرعت چرخش، غلظت H_2O_2 و غلظت نیترات آمونیوم همگی مورد بررسی قرار گرفتند. نتایج، رابطه مطلوبی را بین سرعت انحلال نقره (Ag) و سرعت چرخش نشان می‌دهد. علاوه بر این، غلظت کم نیترات آمونیوم (بین ۰.۲۰ تا ۰.۰۰۳ مولار) اثرات مفیدی بر انحلال نقره دارد. سرعت انحلال به طور قابل توجهی تحت تأثیر غلظت‌های H_2O_2 بین ۰.۰۸ تا ۰.۱۵ مولار قرار گرفت، زیرا این محدوده از غلظت H_2O_2 برای فراهم کردن پتانسیل اکسیداتیو کافی برای حل‌الیت قابل توجه نقره مورد نیاز است. با این حال، این اثر در محدوده ۰.۲۰ تا ۰.۵۰ مولار کمتر مشهود است. محدوده دمایی ۲۰ تا ۵۰ درجه سانتیگراد مفید است زیرا H_2O_2 در این محدوده پایدار است. محاسبه شد که انرژی فعال‌سازی ۲۵.۶۶ کیلوژول بر مول بود.

اطلاعات مقاله

تاریخ ارسال: ۲۰۲۵/۰۴/۰۸

تاریخ داوری: ۲۰۲۵/۰۵/۲۸

تاریخ پذیرش: ۲۰۲۵/۰۹/۱۵

DOI: 10.22044/jme.2025.16031.3089

کلمات کلیدی

نیترات آمونیوم

پراکسید هیدروژن

انحلال

نقره

انرژی فعال‌سازی